



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Asymptomatic congenital tuberculosis: A case report.

Meyer Sauter, Patrick M ; Marques-Maggio, Ewerton ; Relly, Christa ; Keller, Peter M ; Clarenbach, Christian F ; Berger, Christoph

Abstract: RATIONALE: Congenital tuberculosis (TB) is described as a rare, but severe disease. In contrast to the cases with severe symptoms reported so far, we describe a child with asymptomatic congenital TB. **PATIENT CONCERNS:** An 8-week-old girl was investigated because of newly diagnosed TB in her mother, which complained about cough since 21 weeks gestation. Lung biopsy tissue specimens of the mother revealed necrotizing granuloma with a single acid-fast bacillus (AFB) and *Mycobacterium tuberculosis* (MTB) was detected by polymerase chain reaction. Bronchoalveolar lavage was negative for AFB smear and culture, arguing against postnatal transmission of MTB. TB contact investigations were negative. The child, at the age of 8 weeks at first assessment, was in an excellent general condition and diagnosed with congenital TB by culture-positive lung TB and exclusion of postnatal transmission. **DIAGNOSES:** The child fulfilled Cantwell criteria to diagnose congenital TB. **INTERVENTIONS:** Ambulatory anti-tuberculosis treatment was initiated for 6 months. **OUTCOMES:** The 18 months follow-up was uneventful. **LESSONS:** This case of asymptomatic congenital TB in a young child illustrates the diagnostic difficulties in congenital TB and raises the question whether congenital TB is underestimated.

DOI: <https://doi.org/10.1097/MD.00000000000007562>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-138797>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Meyer Sauter, Patrick M; Marques-Maggio, Ewerton; Relly, Christa; Keller, Peter M; Clarenbach, Christian F; Berger, Christoph (2017). Asymptomatic congenital tuberculosis: A case report. *Medicine*, 96(29):e7562.

DOI: <https://doi.org/10.1097/MD.00000000000007562>

Asymptomatic congenital tuberculosis

A case report

Patrick M. Meyer Sauter, MD^{a,b,*}, Ewerton Marques-Maggio, MD^c, Christa Relly, MD^{a,b}, Peter M. Keller, MD^{d,e}, Christian F. Clarenbach, MD^f, Christoph Berger, MD^{a,b}

Abstract

Rationale: Congenital tuberculosis (TB) is described as a rare, but severe disease. In contrast to the cases with severe symptoms reported so far, we describe a child with asymptomatic congenital TB.

Patient concerns: An 8-week-old girl was investigated because of newly diagnosed TB in her mother, which complained about cough since 21 weeks gestation. Lung biopsy tissue specimens of the mother revealed necrotizing granuloma with a single acid-fast bacillus (AFB) and *Mycobacterium tuberculosis* (MTB) was detected by polymerase chain reaction. Bronchoalveolar lavage was negative for AFB smear and culture, arguing against postnatal transmission of MTB. TB contact investigations were negative. The child, at the age of 8 weeks at first assessment, was in an excellent general condition and diagnosed with congenital TB by culture-positive lung TB and exclusion of postnatal transmission.

Diagnoses: The child fulfilled Cantwell criteria to diagnose congenital TB.

Interventions: Ambulatory anti-tuberculosis treatment was initiated for 6 months.

Outcomes: The 18 months follow-up was uneventful.

Lessons: This case of asymptomatic congenital TB in a young child illustrates the diagnostic difficulties in congenital TB and raises the question whether congenital TB is underestimated.

Abbreviations: AFB = acid-fast bacillus, BAL = bronchoalveolar lavage, CT = computed tomography, IGRA = interferon-gamma release assay, IVF = in vitro fertilization, MTB = *Mycobacterium tuberculosis*, PCR = polymerase chain reaction, TB = tuberculosis, TNF = tumor necrosis factor, TST = tuberculin skin test.

Keywords: asymptomatic, congenital, miliary, tree-in-bud opacities, tuberculosis

1. Introduction

Congenital tuberculosis (TB) is difficult to diagnose.^[1] We were recently challenged by a child with culture-proven lung TB acquired by vertical transmission illustrating these diagnostic difficulties in clinical practice. Interestingly, in contrast to the cases of congenital TB with severe symptoms reported so far,^[2] this child did not manifest any symptoms.

Editor: Maria Jose Carbonero Celis.

The authors have no conflicts of interest to disclose.

^a Division of Infectious Diseases and Hospital Epidemiology, ^b Children's Research Center, University Children's Hospital Zurich, ^c Division of Clinical Pathology, University Hospital Zurich, ^d Institute of Medical Microbiology, ^e Swiss National Center for Mycobacteria, University of Zurich, ^f Department of Pulmonology, University Hospital Zurich, Zurich, Switzerland.

* Correspondence: Patrick M. Meyer Sauter, Division of Infectious Diseases and Hospital Epidemiology, University Children's Hospital Zurich, Steinwiesstrasse 75, CH-8032 Zurich, Switzerland (e-mail: patrick.meyer@kispi.uzh.ch).

Copyright © 2017 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Medicine (2017) 96:29(e7562)

Received: 7 June 2017 / Received in final form: 23 June 2017 / Accepted: 28 June 2017

<http://dx.doi.org/10.1097/MD.00000000000007562>

2. Case report

An 8-week-old girl was investigated because of newly diagnosed TB in her mother, which presented the first time 4 weeks after birth with a cough that emerged at 21 weeks gestation, during treatment of indeterminate colitis with the anti-tumor necrosis factor (TNF)- α antibody adalimumab (Fig. 1A). She was HIV negative. No radiographic studies were conducted so far. The interferon-gamma release assay (IGRA), performed before pregnancy under treatment with prednisolone because of colitis, was negative. Her computed tomography (CT) scan 4 weeks after delivery showed numerous nodules and tree-in-bud opacities (Fig. 1A1), a pattern characteristic for miliary TB.^[3] Lung biopsy tissue specimens revealed necrotizing granuloma (Fig. 1A2) with a single acid-fast bacillus (AFB) (Fig. 1A3), and *Mycobacterium tuberculosis* (MTB) was detected by polymerase chain reaction (PCR). Bronchoalveolar lavage (BAL) was negative for MTB, arguing against postnatal transmission of MTB. TB contact investigations led to the investigation of this child and were otherwise negative.

The child, at the age of 8 weeks at first assessment, was in an excellent general condition (Fig. 1B). Pregnancy induced by in vitro fertilization (IVF), birth, and neonatal period were unremarkable. Physical examination, laboratory evaluation, abdominal ultrasound, and chest x-ray were normal (Fig. 1B1). Surprisingly, the tuberculin skin test (TST) was positive with an induration of 9 \times 6 mm at 8 weeks of age, and 20 \times 10 mm at 10 weeks of age. Three gastric aspirates did not show AFB by microscopy, but 1 grew MTB in culture after




Figure 1. The course of TB in the mother (A) and the child (B). There are 2 x axes: the x axis above is the timeline of the medical history of the mother (panel A: January 2013–December 2016), and below the timeline of the medical history of the child (panel B: October 2014–December 2016). Symptoms of the mother are indicated as red areas (x axis above), and pregnancy is shown as orange area (x axis below). Investigations positive for TB are indicated in red (both cultured strains from the mother and the child were identical in sequence and fully susceptible). Treatment regimens are shown as grey arrows. A1: CT scan (axial view) with miliary nodules (circle) and tree-in-bud opacities (arrow). A2: Histopathology of miliary nodules (hematoxylin and eosin stain) showing granuloma characterized by necrotic caseating granuloma center (asterix), epithelioid histiocytes (arrow), and Langhans giant cells (circle). A3: Single acid-fast curvilinear bacillus in lung biopsy tissue specimen (Ziehl-Neelsen stain). B1: Chest x-ray of the child at diagnosis of congenital TB. The TST was performed with 0.1 mL (2 units) PPD RT23 SSI, Statens Serum Institute, Denmark. The placenta was not available for further investigations at diagnosis. Informed consent was given by the mother of the patient. CT= computed tomography; IGRA=interferon-gamma release assay; IVF=in vitro fertilization; PCR=polymerase chain reaction; TST=tuberculin skin test.

3 weeks. Ambulatory anti-tuberculosis treatment was initiated for 6 months (Fig. 1B). The 18 months follow-up was uneventful.

3. Discussion

Both anti-TNF- α treatment and pregnancy are associated with an increased risk of miliary TB.^[3] Infertility (requiring IVF)^[1,4] and

colitis^[3] as present in the mother have been associated with genital and intestinal TB, respectively. In fact, cough and colitis completely resolved after initiation of anti-tuberculosis treatment. Disseminated miliary TB in the immunocompromised mother may be in line with hematogenous spread of MTB to the child in utero.^[3]

Congenital infection is further supported by the strongly positive TST of the child already at 8 weeks of age. TST conversion occurs

within 6 to 8 weeks after infection, and again in congenital TB the TST is reported to convert months after birth.^[5] Together with the negative BAL of the mother 4 weeks after delivery it seems less likely that TB was acquired after birth. We lack the examination of the placenta to eventually proof vertical transmission. Nevertheless, the child indeed fulfilled Cantwell criteria^[5] to diagnose congenital TB, that is, culture-positive lung TB and exclusion of postnatal transmission by thorough contact investigation.

In conclusion, this case illustrates the challenges in diagnosing congenital TB and raises the question whether congenital TB can present as asymptomatic infection.

References

- [1] Shah G, Tse-Chang A, Cooper R, et al. Nodular lung lesions in a 10-week-old infant. *Pediatr Infect Dis J* 2015;34:912, 917.
- [2] Mazade MA, Evans EM, Starke JR, et al. Congenital tuberculosis presenting as sepsis syndrome: case report and review of the literature. *Pediatr Infect Dis J* 2001;20:439–42.
- [3] Sharma SK, Mohan A, Sharma A, et al. Miliary tuberculosis: new insights into an old disease. *Lancet Infect Dis* 2005;5:415–30.
- [4] Flibotte JJ, Lee GE, Buser GL, et al. Infertility, in vitro fertilization and congenital tuberculosis. *J Perinatol* 2013;33:565–8.
- [5] Cantwell MF, Shehab ZM, Costello AM, et al. Brief report: congenital tuberculosis. *N Engl J Med* 1994;330:1051–4.